

PATENT
Docket No. 236472000720

CERTIFICATE OF MAILING BY "FIRST CLASS MAIL"

I hereby certify that this correspondence is being deposited with the United States Postal Service as first class mail in an envelope addressed to:
Assistant Commissioner for Patents, Washington, D.C. 20231, on May 20, 1997.

5-20-97
Date

Jennifer L. Taylor
Jennifer L. Taylor

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In the application of:

Michael C. Kiefer et al.

Serial No.: 08/320,157

Filing Date: October 7, 1994

For: NOVEL APOPTOSIS-MODULATING
PROTEINS, DNA ENCODING THE
PROTEINS AND METHODS OF USE
THEREOF

Examiner: Ray F. Ebert

Group Art Unit: 1806

DECLARATION UNDER 37 C.F.R § 1.132

Assistant Commissioner for Patents
Washington, D.C. 20231

Dear Sir:

I, Michael C. Kiefer, declare as follows:

1. I am an inventor of the above-referenced patent application, and as such am familiar with the claimed subject matter. I have read the Office Action dated December 3, 1996 and understand that the Examiner would like to see evidence that CDN proteins of the claimed invention 1) accumulate intracellularly 2) are in a form that is available and bindable by antibodies made against recombinantly expressed CDN protein and 3) are present in sufficient quantity for detectable binding to antibodies. The following experiments were performed at my direction and show that CDN proteins are present in sufficient quantities and are recognized by

antibodies raised to recombinant CDN protein. A copy of my Curriculum Vitae is attached as Exhibit 1. As shown in my curriculum vitae, I am eminently qualified to supervise these experiments and to interpret the results.

2. Recognition of CDNs by anti-CDN antibody; Experimental protocol.

In these studies, the molecular basis of the effect of interferon- γ (IFN- γ) to increase the sensitivity of some cells to tumor necrosis factor- α - and anti-Fas antibody-mediated cell death was analyzed. A human colon adenocarcinoma cell line, HT-29, was treated with IFN- γ and the expression of various genes and proteins involved in apoptosis was measured. In the experiments described herein, the protein designated Bak is identical to the protein referred to as CDN in the instant invention.

In an attempt to identify genes that may be involved in the increased apoptotic sensitivity of HT-29 cells, the protein levels of various apoptosis modulators were analyzed before and after IFN- γ treatment. Whole cell extracts for Bak, Bax, Bcl-2 and Bcl-x analysis were prepared by lysing 1×10^7 HT-29 cells in 200 μ l of 1X phosphate-buffered saline (PBS) containing 1% NP-40, 0.5% Na-deoxycholate, 0.1% sodium dodecyl sulfate (SDS), 1 mM phenyl methyl sulfonyl fluoride (PMSF), 10 μ g/ml leupeptin and 1 μ g/ml aprotinin for 10 minutes at 4°C. Lysates were centrifuged for 10 minutes at 1000 x g and supernatants were saved. For the protein determinations, lysates from HT-29 cells were quantitated by the Bradford assay. The samples (25-100 μ g protein/lane) were resolved by 15% SDS-polyacrylamide gel electrophoresis (PAGE), transferred to nitrocellulose and incubated for 1 hr in blocking buffer (PBS containing 5% milk powder and 0.05% Tween-20). For Bak analysis, the blot was incubated for 1 hr with rabbit antisera directed against Bak in blocking buffer followed by a 1 hr incubation with anti-rabbit IgG-peroxidase conjugate (1:2000) and then developed using the ECL method (Amersham). The anti-Bak antisera was generated in rabbits using purified recombinant Bak produced in yeast and used at 1:5000 dilution. The anti Bcl-x, Bax and Bcl-2 antibodies were used as recommended by supplier (Santa Cruz Biotechnology, Inc.). The results obtained are depicted in Figure 1.

3. Results.

Northern blot analysis of bcl-2 family members had revealed that bak mRNA levels were induced after 16 hr of IFN- γ treatment, whereas the levels of bax, bcl-x, and bcl-2 mRNA were

unchanged. Bak protein levels also increased in HT-29 cells following IFN- γ treatment as shown by Western blot analysis (Fig. 1). At time zero, Bak protein levels were low but detectable.

The increase in Bak protein reflects the corresponding increase in Bak mRNA levels. Both Bax and Bcl-x protein levels remain unchanged following IFN- γ treatment which is consistent with their constant mRNA levels.




Fig. 1 Western blot analysis of Bak, Bcl-x and Bax protein levels. HT-29 cells were treated with 200 U/ml IFN- γ . At zero, 6 or 16 hours after treatment, cells were lysed and 25-100 μ g total protein was loaded per lane onto a 15% SDS polyacrylamide gel. After electrophoresis, the proteins were transferred to nitrocellulose filters. After blocking the filters, antibodies to either Bak, Bcl-x or Bax were added and allowed to bind. After washing the filter, peroxidase-labelled secondary antibodies were added and, after washing, detected by enhanced chemiluminescence, using a kit (Amersham).

The experiments described herein demonstrate that CDN proteins are intracellular proteins and can be detected in cell lysates. The experiments further demonstrate that antibodies generated using recombinant CDN protein produced in yeast can recognize CDN proteins in cell lysates, but not Bcl-x or Bax. These antibodies therefore recognize CDNs, but not other members of the bcl family tested.

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application, any patent issuing thereon, or any patent to which this verified statement is directed.

4-23-97
Date

Michael C. Kiefer
Michael C. Kiefer

MICHAEL C. KIEFER, Ph.D.

PERSONAL

Home address: 401 Wright Court, Clayton, CA 94517
Home phone: (510) 672-3576
Business address: LXR Biotechnology Inc., 1401 Marina Way South, Richmond, CA 94804
Business phone: (510) 412-9100

EDUCATION

1983 Ph.D. Biochemistry
University California, Davis

1976 Concurrent Undergraduate
University California, Berkeley

1974 B.S. Zoology, Graduated with Honors
University California, Davis

CAREER EXPERIENCE

3/93 to 3/97 LXR Biotechnology Inc.
Richmond, California

2/96 to 3/97 Vice President, Molecular Biology

3/95 to 1/96 Director, Molecular Biology

3/93 to 2/95 Senior Scientist

Responsible for preclinical molecular biological research and development, including strategic decisions, budgeting and research planning in the following areas:

— Gene Discovery: Identification, cloning and characterization of novel apoptosis-related genes and validation as important therapeutic targets.

— Signal Transduction: Identification and characterization of apoptotic pathways involving novel genes and drug leads discovered at LXR.

— Assay Development: Design and development of assays suitable for high-throughput screening of small molecule libraries with the goal of isolating compounds that modulate the apoptotic activity of genes, gene products and other lead drugs discovered at LXR.

Responsible for molecular biology presentations: 1) to pharmaceutical companies and other potential investors; and 2) at international scientific meetings.

Manage and direct a research group of 12 scientists, including 5 Ph.D.s. Research has resulted in 1 issued patent and 5 patents pending.

7/87 to 2/93 **Chiron Corporation**
Emeryville, California

2/91 to 2/93 Senior Scientist

8/87 to 1/91 Research Scientist

Primary responsibility was discovery research in the areas of bone growth and differentiation. Utilized novel and standard cloning techniques to isolate cDNAs encoding new growth factor receptors, bone growth modulators, proteases and malarial antigens. The genes were expressed in yeast, baculovirus or mammalian based expression systems and the functions of the purified recombinant proteins were analyzed in appropriate *in vitro* cell based assays or *in vivo* animal models.

Research resulted in three issued patents and two patents pending.

1/86 to 7/87 **E.I. Du Pont de Nemours & Co., Medical Products & Immunology**
Glenolden, Pennsylvania

Visiting Scientist

Responsible for cloning and expressing genes involved in T and B cell differentiation and function. Primary emphasis was in utilizing cDNA expression vectors and expression systems which, in conjunction with differential cDNA screening, allow identification of the following: 1) Genes that are involved in T cell activation in response to interleukin 2 (IL-2); 2) pre-B cell specific surface antigens; and 3) lymphokines and lymphokine receptors. Other work included characterization and sequencing of two pre-B cell related cDNAs and gene transfer of the IL 2 receptor constructs into fibroblasts for structure/function analysis.

10/83 to 12/85 **Basel Institute for Immunology**
Basel, Switzerland

Staff Scientist

Studied the molecular biology of T cell lymphocytes. Primary emphasis was the isolation and characterization of T cell specific genes including the T cell receptor. Approaches included subtractive cDNA cloning of T cell specific mRNAs. One T cell specific cDNA was further characterized by restriction enzyme analysis, Northern and Southern blot hybridization, DNA sequencing and chromosome mapping. The cDNA was expressed in bacteria as a fusion protein, purified and injected into rabbits to yield antisera that detected the protein in T cells.

10/81 to 9/83 **U.S. Department of Agriculture, Plant Virology Laboratory,**
Plant Protection Institute
Beltsville, Maryland

Postdoctoral Researcher

Investigated the mechanisms of viroid replication and pathogenesis. Approaches used were the following: 1) Comparisons of the primary and secondary structure (derived by computer analysis) of viroids that display a wide range of symptoms in tomato. The sequence of two viroids were determined by direct enzymatic RNA sequencing and chemical sequencing of cDNA clones. 2) Analysis of viroid replication and pathogenesis in tomato, using as inocula, potato spindle tuber viroid and tomato apical stunt viroid cDNA clones, and chimeric and mutant derivatives.

10/76 to 9/81 **University of California, Davis, Dept. of Biochemistry**
Davis, California
Major Professor - Dr. George Bruening

Doctoral Researcher

Investigated the molecular basis of plant resistance to virus infection using a resistant variety of *Vigna unguiculata* (cowpea) protoplasts and cowpea mosaic virus as a model system. Techniques used included liposome mediated delivery of viral RNA into protoplasts, ELISA assay, Northern blotting, *in vivo* radiolabelling of protein and nucleic acids, gel electrophoresis and fluorescent antibody labelling of virus infected protoplasts.

12/75 to 9/76

University of California, Berkeley, Dept. of Biochemistry
Berkeley, California

Research Professor - Dr. Michael Chamberlain

Predoctoral Researcher

Investigated early transcriptional termination in bacteriophages T3 and T7. Techniques used included *in vivo* radiolabelling of virus specific proteins, gel electrophoresis and phage and bacterial selection and growth.

10/74 to 9/75

University of Freiburg, Institut für Biologie III
Freiburg, West Germany
Research Professor - Dr. Gerd Hobom

Predoctoral Researcher

Duties included purification of restriction endonucleases, restriction mapping of bacteriophages, growth and purification of defective bacteriophage lambda and agarose gel electrophoresis.

3/73 to 9/74

University of California, Davis, Dept. of Biological Chemistry
Davis, California
Research Professor - Dr. Robert Traut

Undergraduate Researcher

Investigated the specificity of ATP and GTP dependent protein kinases with respect to their ability to phosphorylate ribosomal proteins of *Escherichia coli*. Techniques used included ribosome purification by zonal centrifugation, two dimensional gel electrophoresis and enzyme assays.

TEACHING EXPERIENCE

9/77 to 7/80

University of California, Davis, Dept. of Biochemistry and Biophysics
Davis, California

Teaching Assistant

Led discussion sections in undergraduate biochemistry laboratory and lecture courses.

PUBLICATIONS

1. Issinger, O.G., Kiefer, M.C., and Traut, R.R. (1975). Specificity of ATP-dependent and GTP-dependent protein kinases with respect to ribosomal proteins of *Escherichia coli*. *Eur. J. Biochem.* 59, 137-143.
2. Kiefer, M.C., Neff, N., and Chamberlain, M.J. (1977). Transcriptional termination at the end of the early region of bacteriophages T3 and T7 is not affected by polarity suppressors. *J. Virol.* 22, 548-552.
3. Bruening, G., and Kiefer, M. (1981). Control of immunity to cowpea mosaic virus. In *Proceedings of Symposia, IX International Congress of Plant Protection*, 1, 225-228.
4. Kiefer, M.C., Daubert, S.D., Schneider, I.R., and Bruening, G. (1982). Multimeric forms of satellite of tobacco ringspot virus RNA. *Virology* 121, 262-273.
5. Cress, D.E., Kiefer, M.C., and Owens, R.A. (1983). Biological activity of cloned potato spindle tuber viroid cDNA. In *Plant Infectious Agents - Viruses, Viroids, Virusoids and Satellites*, H.D. Robertson, S.H. Howell, M. Zaitlin, R.L. Malmberg, eds. (New York: Cold Spring Harbor Laboratory), 160-164.
6. Cress, D.E., Kiefer, M.C., and Owens, R.A. (1983). Construction of infectious potato spindle tuber viroid cDNA clones. *Nucl. Acids. Res.* 11, 6821-6835.
7. Eastwell, K.C., Kiefer, M.C., and Bruening, G. (1983). Immunity of cowpeas to cowpea mosaic virus. In *Plant Molecular Biology, ARCO-UCLA Symposia on Molecular and Cell Biology*, R. Goldberg, ed. (New York: Academic Press), XII, 201-211.
8. Kiefer, M.C., Bruening, G., and Russell, M.L. (1983). RNA and nucleocapsid accumulation in protoplasts resistant to cowpea mosaic virus. In *Plant Infectious Agents - Viruses, Viroids, Virusoids and Satellites*, H.D. Robertson, S.H. Howell, M. Zaitlin, R.L. Malmberg, eds. (New York: Cold Spring Harbor Laboratory), 106-110.
9. Kiefer, M.C., Owens, R.A., and Diener, T.O. (1983). Structural similarities between viroids and transposable genetic elements. *Proc. Natl. Acad. Sci. USA* 80, 6234-6238.
10. Kiefer, M.C., Bruening, G., and Russell, M.L. (1984). RNA and capsid accumulation in cowpea protoplasts that are resistant to cowpea mosaic virus strain SB. *Virology* 137, 371-381.
11. Goridis, C., Hirn, M., Santoni, M.-J., Gennarini, G., Deagostini-Bazin, H., Jordan, B. R., Kiefer, M., and Steinmetz, M. (1985). Isolation of mouse N-CAM-related cDNA: detection and cloning using monoclonal antibodies. *EMBO J.* 4, 631-635.

12. Hammond, R.W., Kiefer, M.C., Cress, D.E., and Owens, R.A. (1985). Probing viroid structure-function relationships by site-specific mutagenesis of infectious cDNA. In *Molecular Form and Function of the Plant Genome*, L. van Vloten-Doting, G.S.P. Groot, T.C. Hall, eds. (New York: Plenum Publishing), 387-398.
13. Nemazee, D.A., Studer, S., Steinmetz, M., Dembic, Z., and Kiefer, M. (1985). The lymphoproliferating cells of MRL-*lpr/lpr* mice are a polyclonal population that bear the T lymphocyte receptor for antigen. *Eur. J. Immunol.* *15*, 760-764.
14. Owens, R.A., and Kiefer, M.C. (1985). Cloning of viroid cDNA. In *Recombinant DNA Research and Virus*, Y. Becker, ed. (Boston: Nijhoff), 293-304.
15. Snodgrass, H.R., Kisielow, P., Kiefer, M., Steinmetz, M., and von Boehmer, H. (1985). Ontogeny of the T cell antigen receptor within the thymus. *Nature* *313*, 592-595.
16. Huppi, K., D'Hoostelaere, L., Kiefer, M., Steinmetz, M., and Jouvin-Marche, E. (1986). The context of T-cell receptor gamma chain genes among wild mouse species. *Immunogenetics* *24*, 304-308.
17. Owens, R.A., Hammond, R.W., Gardner, R.C., Kiefer, M.C., Thompson, S.M., and Cress, D.E. (1986). Site-specific mutagenesis of potato spindle tuber viroid cDNA. *Plant Mol. Biol.* *6*, 197-192.
18. Traunecker, A., Kiefer, M., Dembic, Z., Steinmetz, M., and Karjalainen, K. (1986). Rearrangements of T cell receptor loci can be found only rarely in B lymphoid cells. *Eur. J. Immunol.* *16*, 430-434.
19. Neeper, M.P., Kuo, L.-M., Kiefer, M.C., and Robb, R.J. (1987). Structure-function relationships for the IL 2 receptor system III. Tac protein missing amino acids 102-173 (Exon 4) is unable to bind IL 2. Detection of spliced protein after L cell transfection. *J. Immunol.* *138*, 3532-3538.
20. Palacios, R., Kiefer, M., Brockhaus, M., Karjalainen, K., Dembic, Z., Kisielow, P., and von Boehmer, H. (1987). Molecular, cellular and functional properties of bone marrow T lymphocyte progenitor clones. *J. Exp. Med.* *166*, 12-32.
21. Kiefer, M.C., Bauer, D.M., Young, D., Hermsen, K.M., Maslarz, F.R., Barr, P.J. (1988). The cDNA and derived amino acid sequences for human and bovine matrix Gla protein. *Nucl. Acids Res.* *16*, 5213.
22. Kiefer, M.C., Bauer, D.M., Barr, P.J. (1989). The cDNA and derived amino acid sequence for human osteopontin. *Nucl. Acids Res.* *17*, 3306.

23. Power, M.D., Kiefer, M.C., Barr, P.J., Reeves, R. (1989). Nucleotide sequence of human mitochondrial cytochrome c oxidase II cDNA. *Nucl. Acids Res.* 17, 6734.
24. Richards, S., Bucan, M., Brorson, K., Kiefer, M.C., Hunt, S.W. I., Lehrach, H., and Fischer-Lindahl, K. (1989). Genetic and molecular mapping of the Hmt region of mouse. *EMBO J.* 8, 3749-3757.
25. Barr, P.J., Mason, O.B., Wong, P.A., Gibson, H.L., Kiefer, M.C., and Brake, A.J. (1990). Yeast expression: secretion and intracellular systems. In *Recombinant Systems in Protein Expression*, K.K. Alitalo, M.-. Huhtala, J. Knowles, A. Vaheri, eds. Elsevier Science Publishers B.V.), 37-46.
26. Kiefer, M.C., Saphire, A.C.S., Bauer, D.M., Barr, P.J. (1990). The cDNA and derived amino acid sequences of human and bovine bone Gla protein. *Nucl. Acids Res.* 18, 1909.
27. Kiefer, M.C., Stephans, J.C., Crawford, K., Okino, K., and Barr, P.J. (1990). Ligand-affinity cloning and structure of a cell surface heparan sulfate proteoglycan that binds basic fibroblast growth factor. *Proc. Natl. Acad. Sci. USA* 87, 6985-6989.
28. Reeves, R., Spies, G.A., Kiefer, M., Barr, P.J., and Power, M. (1990). Primary structure of the putative human oncogene, *pim-1*. *Gene* 90, 303-307.
29. Schild, D., Brake, A.J., Kiefer, M.C., Young, D., and Barr, P.J. (1990). Cloning of three human multifunctional de novo purine biosynthetic genes by functional complementation of yeast mutations. *Proc. Natl. Acad. Sci.* 87, 2916-2920.
30. Zapf, J., Kiefer, M., Merryweather, J., Masiarz, F., Bauer, D., Born, W., Fischer, J.A., and Froesch, E. R. (1990). Isolation from adult human serum of four insulin-like growth factor (IGF) binding proteins and molecular cloning of one of them that is increased by IGF I administration and in extrapancreatic tumor hypoglycemia. *J. Biol. Chem.* 265, 14892-14898.
31. Wise, R.J., Barr, P.J., Wong, P.A., Kiefer, M.C., Brake, A.J., and Kaufman, R.J. (1990). Expression of a human proprotein processing enzyme: correct cleavage of the von Willebrand factor precursor at a paired basic amino acid site. *Proc. Natl. Acad. Sci. USA* 87, 9378-9382.
32. Barr, P.J., Mason, O.B., Wong, P.A., Lee Ng, C.T., Kiefer, M.C., and Brake, A.J. (1991). Heterologous expression in *Saccharomyces cerevisiae* and other yeasts. *Proceedings of the 6th International Symposium on the Genetics of Industrial Microorganisms*, H. Heslot, J. Davies, J. Florent, L. Bobichon, G. Durand, L. Penasse, eds. (Société Française de Microbiologie), pp. 95-105.

33. Kiefer, M.C., Masiarz, F.R., Bauer, D.M., and Zapf, J. (1991). Identification and molecular cloning of two new 30 kD insulin-like growth factor binding proteins isolated from adult human serum. *J. Biol. Chem.* 266, 9043-9049.
34. Barr, P.J., Mason, O.B., Landsberg, K.E., Wong, P.A., Kiefer, M.C., and Brake, A.J. (1991). cDNA and gene structure for a human subtilisin-like protease with cleavage specificity for paired basic amino acid residues. *DNA and Cell Biol.* 10, 319-328.
35. Kiefer, M.C. Baird, A., Nguyen, T., George-Nascimento, C., Mason, O.B., Boley, L.J., Stephans, J.C.; Valenzuela, P.V., and Barr, P.J. (1991). Molecular cloning of a human basic fibroblast growth factor receptor cDNA and expression of a biologically active extracellular domain in a baculovirus system. *Growth Factors* 5, 115-127.
36. Kiefer, M.C., Joh, R.S., Bauer, D.M., and Zapf, J. (1991). Molecular cloning of a new human insulin-like growth factor binding protein. *Biochem. and Biophys. Res. Comm.* 176, 219-225.
37. Kiefer, M.C., Tucker, J.E., Joh, R., Landsberg, K.E., Saltman, D., and Barr, P.J. (1991). Identification of a second human subtilisin-like protease gene in the *fes/fps* region of chromosome 15. *DNA and Cell Biol.* 10, 757-769.
38. Zapf, J., Schmid, C., Kiefer, M., Guler, H.-P., and Froesch, E.R. (1991). *In vivo* models of IGF binding protein (IGFBP) regulation with possible relevance to their physiological roles. In *Modern Concepts of Insulin-Like Growth Factors, Proceedings of the Second International Symposium on Insulin-Like Growth Factors*, E.M. Spencer, ed. (New York, Elsevier), 591-605.
39. Kiefer, M.C., Ishihara, M., Swiedler, S.J., Crawford, K., Stephans, J.C., and Barr, P.J. (1991). The molecular biology of heparan sulfate fibroblast growth factors receptors. *New York Acad. Sci.* 638, 167-176.
40. Gibson, H.L., Tucker, J.E., Kaslow, D.C., Krettli, A.U., Collins, W.E., Kiefer, M.C., Bathurst, I.C., Barr, P.J. (1992). Structure and expression of the gene for Pv200, a major blood-stage surface antigen of *Plasmodium vivax*. *Mol. Biochem. Parasitol.* 50, 325-334.
41. Ishihara, M., Tyrrell, D.J., Kiefer, M.C., Barr, P.J., and Swiedler, S.J. (1992). A cell-based assay to evaluate the interaction of heparin-like molecules and basic fibroblast growth factor, *Anal. Biochem.* 202, 310-315.
42. Kiefer, M.C., Schmid, C., Waldvogel, M., Schläpfer, I., Futo, E., Green, K.M., Masiarz, F. R., Barr, P.J., and Zapf, J. (1992). Characterization of recombinant insulin-like growth factor binding proteins -4, -5 and -6 produced in yeast. *J. Biol. Chem.* 267, 12692-12699.

43. Ishihara, M., Kiefer, M.C., Barr, P.J., Guo, Y., and Swiedler, S.J. (1992). Selection of COS cell mutants defective in the biosynthesis of heparan sulfate proteoglycan. *Anal. Biochem.* 206, 400-407.
44. Conover, C.A., Kiefer, M.C., and Zapf, J. (1993). Post-translational regulation of insulin-like growth factor (IGF) binding protein-4 in normal and transformed human fibroblasts: IGF dependence and biological studies. *J. Clin. Invest.* 91, 1129-1137.
45. Tyrrell, D.J., Ishihara, M., Rao, N., Horne, A., Kiefer, M.C., Stauber, G.B., Lam, L.H., and Stack, R.J. (1993). The structure and biological activities of a heparin-derived hexasaccharide with high affinity for basic fibroblast growth factor. *J. Biol. Chem.* 268, 4684-4689.
46. Conover, C.A. and Kiefer, M.C. (1993). Regulation and biological effect of endogenous insulin-like growth factor binding protein-5 in human osteoblastic cells. *J. Clin. Endo. Metab.* 76, 1153-1159.
47. Rohan, R.M., Ricciarelli, E., Kiefer, M.C., Resnick, C.E., and Adashi, E.Y. (1993). Rat ovarian insulin-like growth factor binding protein-6: A pituitary-dependent theca-interstitial selective species with limited antigonadotropic activity. *Endocrinol.* 132, 2507-2512.
48. Andress, D.L., Loop, S.M., Boley, L.J., and Kiefer, M.C. (1993). Mitogenic activity of truncated insulin-like growth factor (IGF) binding protein-5 in osteoblast-like cells. *Biochem. and Biophys. Res. Comm.* 195, 25-30.
49. Kiefer, M.C., Schmid, C., Waldvogel, M., Schläpfer, I., Futo, E., Masiarz, F. R., Green, K., Barr, P.J., and Zapf, J. (1993). Recombinant human insulin-like growth factor binding proteins 4, 5, and 6: biological and physicochemical characterization. *Growth Reg.* 3, 56-59.
50. Durham, S.K., Kiefer, M.C., Riggs, B.L., and Conover, C.A. (1994). Evidence for a specific insulin-like growth factor (IGF)-dependent IGF binding protein-4 protease in normal human osteoblast-like cells. *J. Bone and Min. Res.* 1, 111-117.
51. Caroni, P., Schneider, C., Kiefer, M.C., and Zapf, J. (1994). Role of muscle insulin-like growth factors in nerve sprouting: suppression of terminal sprouting in paralyzed muscle by IGF-binding protein 4. *J. Cell Biol.* 125, 893-902.
52. Allander, S.V., Larsson, C., Ehrenborg, E., Suwanichkul, A., Weber, G., Morris, S.L., Bajalica, S., Kiefer, M.C., Luthman, H., and Powell, D.R. (1994). Characterization of the chromosomal gene and promoter for human insulin-like growth factor binding protein-5. *J. Biol. Chem.* 269, 10891-10898.

53. Cheng, J., Zhou, T., Liu, C., Shapiro, J.P., Brauer, M.J., Kiefer, M.C., Barr, P.J., and Mountz, J.D. (1994). Protection from Fas-mediated apoptosis by a soluble form of the Fas molecule. *Science* 263, 1759-1762.
54. Conover, C.A., Durham, S.K., Zapf, J., Masiarz, F.R., and Kiefer, M.C. (1995). Cleavage analysis of insulin-like growth factor (IGF)-dependent IGF-binding protein-4 proteolysis and expression of protease resistant IGF-binding protein-4 mutants. *J. Biol. Chem.* 270, 4395-4400.
55. Booth, B.A., Boes, M., Andress, D.L., Dake, B.L., Kiefer, M.C., Maack, C., Linhardt, R.J., Bar, K., Caldwell, E.E.O., Weiler, J., and Bar, R.S. (1995). IGFBP-3 and IGFBP-5 Association with Endothelial Cells: Role of C-terminal Heparin Binding Domain. *Growth Reg.* 5, 1-17.
56. Pavloff, N., Potempa, J., Pike, R.N., Prochazka, V., Kiefer, M.C., Travis, J., and Barr, P.J. (1995). Molecular cloning and structural characterization of the Arggingipain proteinase of *Porphyromonas gingivalis*. Biosynthesis as a proteinase-adhesin polyprotein. *J. Biol. Chem.* 270, 1007-1010.
57. Pemberton, P.A., Wong, D.T., Gibson, H.L., Kiefer, M.C., Fitzpatrick, P.A., Sager, R., and Barr, P.J. (1995). The tumor suppressor maspin does not undergo the stressed to relaxed transition or inhibit trypsin-like serine proteases. Evidence that maspin is not a protease inhibitory serpin. *J. Biol. Chem.* 270, 15832-15837.
58. Kiefer, M.C., Brauer, M.J., Powers, V.C., Wu, J.J., Umansky, S.R., Tomei, L.D., and Barr, P.J. (1995). Modulation of apoptosis by the widely distributed Bcl-2 homologue Bak. *Nature* 374, 736-739.
59. Kiefer, M.C., Crawford, K.A., Boley, L.J., Landsberg, K.E., Gibson, H.L., Kaslow, D.C., and Barr, P.J. (1996). Identification and cloning of a locus of serine repeat antigen (sera)-related genes from *Plasmodium vivax*. *Mol. Biochem. Parasitol.* 78, 55-65.
60. Guo, Z., Liliom, K., Fischer, D.J., Bathurst, I.C., Tomei, L.D., Kiefer, M.C., and Tigyi, G. (1996). Molecular cloning of a high-affinity receptor for the growth factor-like lipid mediator lysophosphatidic acid from *Xenopus* oocytes. *Proc. Natl. Acad. Sci. USA* 93, 14367-14372.
61. Pavloff, N., Pemberton, P.A., Potempa, J., Chen, W.C., Pike, R.N., Prochazka, V., Kiefer, M.C., Travis, J., and Barr, P.J. (1997). Molecular cloning and characterization of *Porphyromonas gingivalis* lysine-specific gingipain. A new member of an emerging family of pathogenic bacterial cysteine proteinases. *J. Biol. Chem.* 272, 1595-1600.

62. Pemberton, P.A., Tipton, A.R., Pavloff, N., Smith, J., Erickson, J.R., Mouchabeck, Z.M., Shih, M., Tomei, L.D., Barr, P.J., and Kiefer, M.C. (1997). Tissue distribution and subcellular localization of maspin. Implications for its function as a tumor suppressor. (manuscript submitted).
63. Ossina, N.K., Cannas, A., Powers, V.C., Fitzpatrick, P.A., Knight, J.D., Gilbert, J.R., Shekhtman, E.M., Tomei, L.D., Umansky, S.R., and Kiefer, M.C. (1997).

Interferon- γ modulates a p53-independent apoptotic pathway and apoptosis-related gene-expression. (manuscript submitted).

INVITED SPEAKER

American Chemical Society Meeting, San Francisco, April 1992. "Characterization of recombinant insulin-like growth factor binding proteins -4, -5, and -6 produced intracellularly in yeast."

Ninth International Congress on Endocrinology, Workshop on IGF binding proteins, Opio, France, August 1992. "Characterization of recombinant insulin-like growth factor binding proteins -4, -5, and -6 produced intracellularly in yeast."

American Society for Biochemistry and Molecular Biology Fall Symposium, Genetic and Biochemical Approaches for Studying Cell Death, Granlibakken, Lake Tahoe, California, October 1994. "Cdn-1, -2, and -3, a family of Bcl-2-related genes that modulate apoptosis."

Fourth Abano Terme Meeting on Rehabilitation, First International Conference on Apoptosis in Skeletal and Cardiac Muscles (ASCM96), Abano Terme (Padova), Italy, June 1996. "Interferon- γ sensitizes cells to apoptotic stimuli and increases apoptosis-related gene expression."

REFERENCES

References will be furnished upon request.